

## PRIOR AUTHORIZATION POLICY

**POLICY:** Oncology – Gilotrif Prior Authorization Policy

- Gilotrif<sup>™</sup> (afatinib tablets – Boehringer Ingelheim)

**REVIEW DATE:** 12/01/2021; selected revision 01/19/2022

---

### OVERVIEW

Gilotrif, a tyrosine kinase inhibitor (TKI), is indicated for the following uses:<sup>1</sup>

- **Non-small cell lung cancer (NSCLC)**, first-line treatment of patients with metastatic disease whose tumors have non-resistant epidermal growth factor receptor (*EGFR*) mutations as detected by an FDA-approved test. Limitations of use: The safety and efficacy of Gilotrif have not been established in patients whose tumors have resistant *EGFR* mutations.
- **NSCLC, squamous cell**, for the treatment of patients with metastatic disease progressing after platinum-based chemotherapy.

### Guidelines

Gilotrif has been addressed in National Comprehensive Cancer Network (NCCN) guidelines.<sup>2-4</sup>

- **Head and Neck Cancer:** Guidelines (version 1.2022 – December 8, 2021) recommend Gilotrif as a single agent for the treatment of recurrent, unresectable, or metastatic non-nasopharyngeal cancers (lip, oral cavity, oropharynx, hypopharynx, glottis, larynx, supraglottic, larynx, ethmoid sinus, maxillary sinus, occult primary) in patients with disease progression or after platinum-based therapy.<sup>3</sup>
- **Non-Small Cell Lung Cancer (NSCLC):** Guidelines (version 1.2022 – December 7, 2021) recommend testing for sensitizing EGFR mutations in patients with metastatic disease.<sup>4</sup> Patients with sensitizing EGFR mutations have a significantly better response to the EGFR tyrosine kinase inhibitors (TKIs) [erlotinib, Gilotrif, Iressa, Tagrisso, and Vizimpro]. The most common EGFR mutations are exon 19 deletions and exon 21 (L858R) substitution mutations. Other less common mutations that are also sensitive to EGFR TKIs include L861Q, G719X, and S768I; these mutations cumulatively account for approximately 10% of all EGFR mutations. NCCN recommends the EGFR TKIs as first-line treatment for patients with advanced or metastatic NSCLC with EGFR exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X, and S768I. NCCN does not recommend Gilotrif for use as second-line treatment for patients with squamous cell NSCLC; NCCN notes Gilotrif to be less efficacious and safe compared with other available options.

### POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Gilotrif. All approvals are provided for 3 years in duration as noted below.

**Automation:** None.

---

## RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Gilotrif is recommended in those who meet the following criteria:

### FDA-Approved Indications

- 1. Non-Small Cell Lung Cancer – Epidermal Growth Factor Receptor (*EGFR*) Mutation-Positive.** Approve for 3 years if the patient meets the following criteria (A, B, and C):
  - A) Patient is  $\geq 18$  years of age; AND
  - B) Patient has advanced or metastatic disease; AND
  - C) Patient has sensitizing *EGFR* mutation-positive non-small cell lung cancer as detected by an approved test.  
Note: Examples of sensitizing *EGFR* mutation-positive non-small cell lung cancer include the following: exon 19 deletions, exon 21 (L858R) substitution mutations, L861Q, G719X, and S768I.
- 2. Non-Small Cell Lung Cancer – Squamous Cell Carcinoma.** Approve for 3 years if the patient meets the following criteria (A, B, and C):
  - A) Patient is  $\geq 18$  years of age; AND
  - B) Patient has metastatic squamous cell carcinoma; AND
  - C) Patient has disease progression after treatment with platinum-based chemotherapy.

### Other Uses with Supportive Evidence

- 3. Head and Neck Cancer.** Approve for 3 years if the patient meets the following criteria (A, B, and C):
  - A) Patient is  $\geq 18$  years of age; AND
  - B) Patient has non-nasopharyngeal head and neck cancer; AND  
Note: Examples of non-nasopharyngeal head and neck cancer are lip, oral cavity, oropharynx, hypopharynx, glottis, larynx, supraglottic larynx, ethmoid sinus, maxillary sinus, occult primary.
  - C) Patient has disease progression on or after platinum-based chemotherapy.

## CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Gilotrif is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

## REFERENCES

1. Gilotrif<sup>™</sup> tablets [prescribing information]. Ridgefield, CT: Boehringer Ingelheim; October 2019.
  2. The NCCN Drugs & Biologics Compendium. © 2022 National Comprehensive Cancer Network, Inc. Available at: <http://www.nccn.org>. Accessed on January 12, 2022. Search terms: afatinib.
  3. The NCCN Head and Neck Cancer Clinical Practice Guidelines in Oncology (version 1.2022 – December 8, 2021). © 2021 National Comprehensive Cancer Network, Inc. Available at: <http://www.nccn.org>. Accessed on January 12, 2022.
  4. The NCCN Non-Small Cell Lung Cancer Clinical Practice Guidelines in Oncology (version 1.2022 – December 7, 2021). © 2021 National Comprehensive Cancer Network, Inc. Available at: <http://www.nccn.org>. Accessed on January 12, 2022.
-