

PRIOR AUTHORIZATION POLICY

POLICY: Oncology – Tagrisso Prior Authorization Policy

- Tagrisso® (osimertinib tablets – AstraZeneca)

REVIEW DATE: 01/31/2024; selected revision 10/16/2024

OVERVIEW

Tagrisso, a tyrosine kinase inhibitor, is indicated for the following uses:¹

- **Non-Small Cell Lung Cancer (NSCLC) – Epidermal growth factor receptor (*EGFR*) Mutation-Positive:**
 - First-line treatment of **metastatic NSCLC** tumors that have *EGFR* exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test, in adults.
 - Tagrisso, **in combination with Alimta** (pemetrexed for intravenous use) **and platinum-based chemotherapy** is indicated for the first-line treatment of locally advanced or metastatic NSCLC that have *EGFR* exon 19 or exon 21 L858R mutations, as detected by an FDA-approved test, in adults.
- **NSCLC – *EGFR* T790M Mutation-Positive:** Treatment of metastatic *EGFR* T790M mutation-positive NSCLC, as detected by an FDA-approved test, in adults whose disease has progressed on or after *EGFR* tyrosine kinase inhibitor (TKI) therapy.
- **NSCLC – *EGFR* Mutation-Positive, Post Tumor Resection:** Adjuvant therapy after tumor resection in adults with NSCLC whose tumors have *EGFR* exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test.
- **NSCLC – *EGFR* Mutation-Positive, Unresectable (Stage III) Disease:** Treatment of locally advanced, unresectable (stage III) NSCLC in adults whose disease has not progressed during or following concurrent or sequential platinum-based chemoradiation therapy and whose tumors have *EGFR* exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test.

Guidelines

National Comprehensive Cancer Network (NCCN) guidelines for NSCLC (version 10.2024 – September 23, 2024) recommend testing for *EGFR* mutations in patients with metastatic disease.² The most common *EGFR* mutations are exon 19 deletion and exon 21 (L858R) substitution mutations. Other less common mutations that are also responsive to *EGFR* tyrosine kinase inhibitors (TKIs) include L861Q, G719X, and S768I. NCCN recommends Tagrisso as the “Preferred” first-line treatment for patients with *EGFR* exon 19 deletion or exon 21 (L858R) substitution mutations. Tagrisso can also be used in combination with Alimta® (pemetrexed for injection) and either cisplatin or carboplatin in the first-line setting (category 1, “Other Recommended” regimens). Tagrisso is also a recommended “Preferred” first-line therapy (category 2A) for *EGFR* mutations L861Q, G719X, and S768I. Tagrisso is also recommended as subsequent treatment for all of these mutations. The panel recommends T790M (a secondary mutation in *EGFR*) testing in patients who progress on erlotinib tablets, Gilotrif® (afatinib tablets), Iressa® (gefitinib tablets), or Vizimpro® (dacomitinib tablets). If the patient has *EGFR* T790M-positive metastatic NSCLC, Tagrisso is recommended as subsequent therapy (category 1). If the disease is *EGFR* T790M-negative, the patient can be continued on the current TKI (i.e., erlotinib, Gilotrif, Iressa, or Vizimpro). Tagrisso is also recommended for use in patients with completely resected stage IB-III A *EGFR* (exon 19 deletion, L858R) NSCLC who received previous adjuvant chemotherapy or are ineligible to receive platinum-based chemotherapy. Tagrisso is recommended for stage IIIA unresectable disease (category 1) after definitive chemoradiation for *EGFR* exon 19 deletion or L858R mutation. A footnote also states that for patients who have received sequential chemoradiation, Imfinzi (durvalumab intravenous infusion) can be considered, or if *EGFR* exon 19 deletion or L858R, Tagrisso is recommended.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Tagrisso. All approvals are provided for the duration noted below.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Tagrisso is recommended in those who meet one of the following criteria:

FDA-Approved Indications

1. Non-Small Cell Lung Cancer. Approve for 1 year if the patient meets ALL of the following (A, B and C):

A) Patient is \geq 18 years of age; AND

B) Patient has advanced or metastatic disease; AND

C) Patient meets ONE of the following (i or ii):

i. Patient has epidermal growth factor receptor (*EGFR*) mutation-positive disease as detected by an approved test; OR

Note: Examples of *EGFR* mutation-positive non-small cell lung cancer include the following: exon 19 deletion, exon 21 (L858R) substitution mutations, L861Q, G719X, and S768I.

ii. Patient meets BOTH of the following (a and b):

a) Patient has epidermal growth factor receptor (*EGFR*) T790M mutation-positive disease as detected by an approved test; AND

b) Patient has progressed on treatment with at least one of the *EGFR* tyrosine kinase inhibitors.

Note: *EGFR* tyrosine kinase inhibitors are erlotinib, Iressa (gefitinib tablets), Vizimpro (dacomitinib tablets), Gilotrif (afatinib tablets).

2. Non-Small Cell Lung Cancer – Post Tumor Resection. Approve for 1 year if the patient meets ALL of the following (A, B, C, and D):

A) Patient is \geq 18 years of age; AND

B) Patient has completely resected disease; AND

C) Patient has *EGFR* exon 19 deletion or exon 21 (L858R) substitution mutation as detected by an approved test; AND

D) Patient meets ONE of the following (i or ii):

i. Patient received previous adjuvant chemotherapy; OR

ii. Patient is ineligible to receive platinum-based chemotherapy.

3. Non-Small Cell Lung Cancer – Unresectable, Stage III. Approve for 1 year if the patient meets ALL of the following (A, B, C, and D):

A) Patient is \geq 18 years of age; AND

B) Patient has locally advanced, unresectable (stage III) disease; AND

C) Patient has *EGFR* exon 19 deletions or exon 21 (L858R) substitution mutation as detected by an approved test; AND

D) Patient has not had disease progression during or following platinum-based chemoradiation therapy.

Note: Patient could have received concurrent or sequential chemoradiation therapy.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Tagrisso is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Tagrisso™ tablets [prescribing information]. Wilmington, DE: AstraZeneca; September 2024
2. The NCCN Non-Small Cell Lung Cancer Clinical Practice Guidelines in Oncology (version 10.2024 – September 23, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on October 14, 2024.

HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	<p>Non-Small Cell Lung Cancer: Criteria for Epidermal Growth Factor Receptor (<i>EGFR</i>) Mutation-Positive (Other than <i>EGFR</i> T790M-positive mutation) AND Epidermal Growth Factor (<i>EGFR</i>) T790M Mutation-Positive are combined into one set of criteria. There was one change to the criterion: previously for T790 mutation-positive disease, the criterion approved if patient has metastatic disease, now, the criterion will approve if the patient has advanced or metastatic disease.</p> <p>Non-Small Cell Lung Cancer- Post Tumor Resection: Added the word “tumor” to criteria set; previously it read “Post Resection”.</p>	01/11/2023
Annual Revision	<p>Non-Small Cell Lung Cancer: Deleted the word “sensitizing” in reference to <i>EGFR</i> mutations. This verbiage is no longer used in the guidelines.</p>	01/31/2024
Update	<p>03/11/2024: No criteria changes. Updated Overview section with new indication for Tagrisso. Also updated Guidelines section.</p>	NA
Selected Revision	<p>Non-Small Cell Lung Cancer – Unresectable, Stage III. Added new approval condition and criteria based on new indication.</p>	10/16/2024