

PRIOR AUTHORIZATION POLICY

POLICY: Lupus – Benlysta Intravenous Prior Authorization Policy

- Benlysta® (belimumab intravenous infusion – GlaxoSmithKline)

REVIEW DATE: 03/19/2025

OVERVIEW

Benlysta intravenous, a B-lymphocyte stimulator (BLyS)-specific inhibitor, is indicated for the following uses:¹

- **Lupus nephritis**, in patients ≥ 5 years of age with active disease who are receiving standard therapy.
- **Systemic lupus erythematosus (SLE)**, in patients ≥ 5 years of age with active disease who are receiving standard therapy.

Limitations of Use: Benlysta has not been studied and is not recommended in patients with severe active central nervous system lupus.¹ The efficacy of Benlysta for the treatment of SLE was studied in patients with a history of autoantibodies (anti-nuclear antibody and/or anti-double-stranded DNA) and an exploratory analysis of the pivotal trial indicated Benlysta was beneficial in patients who were autoantibody positive.

Of note, subcutaneous Benlysta is not targeted in this policy.

Guidelines

Lupus Nephritis

Guidelines for the management of lupus nephritis from Kidney Disease: Improving Global Outcomes (KDIGO) [2024] recommendations include Benlysta or Lupkynis in combination with other medications plus glucocorticoids as initial treatment options for patients with active Class III or IV biopsy confirmed lupus nephritis (strong recommendation, moderate certainty of evidence).³ No preference is given between the treatment protocol options; however, the KDIGO guidelines do provide individual patient clinical factors to consider, including but not limited to, kidney function and histology, risk of disease flare, proteinuria, background suppression, and need for parenteral therapy.

SLE

European League Against Rheumatism (EULAR) guidelines for SLE (2023 update) recommend hydroxychloroquine for all patients, unless contraindicated.² Depending on the type and severity of organ involvement, glucocorticoids can be used but dosing should be minimized or withdrawn when possible. Methotrexate, azathioprine, mycophenolate, and/or biologic agents (Benlysta, Saphnelo® [anifrolumab-fnia intravenous infusion]) should be considered in patients who do not respond to hydroxychloroquine $\pm$ glucocorticoids. EULAR also states biologic agents (Benlysta, Saphnelo) should be considered as second-line therapy for the treatment of patients with active skin disease. Patient with active proliferative lupus nephritis should also consider combination therapy with biologic agents (Benlysta, Lupkynis™ [voclosporin capsules]). In general, the pharmacological interventions should be directed by patient specific characteristics and the type/severity of organ involvement.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Benlysta intravenous. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with Benlysta intravenous as well as the monitoring required for adverse events and long-term efficacy, approval requires Benlysta intravenous to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Benlysta intravenous is recommended in those who meet one of the following:

FDA-Approved Indications

1. **Lupus Nephritis.** Approve for the duration noted if the patient meets ONE of the following (A or B):
 - A) **Initial Therapy.** Approve for 6 months if the patient meets ALL of the following (i, ii, iii, and iv):
 - i. Patient is ≥ 5 years of age; AND
 - ii. Diagnosis of lupus nephritis has been confirmed on biopsy; AND
Note: For example, World Health Organization class III, IV, or V lupus nephritis.
 - iii. The medication is being used concurrently with an immunosuppressive regimen; AND
Note: Examples of an immunosuppressive regimen include azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate mofetil, and/or a systemic corticosteroid.
 - iv. The medication is prescribed by or in consultation with a nephrologist or rheumatologist; OR
 - B) **Patient is Currently Receiving Benlysta Intravenous or Subcutaneous.** Approve for 1 year if the patient meets ALL of the following (i, ii, and iii):
 - i. The medication is being used concurrently with an immunosuppressive regimen; AND
Note: Examples of an immunosuppressive regimen include azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate mofetil, and/or a systemic corticosteroid.
 - ii. The medication is prescribed by or in consultation with a nephrologist or rheumatologist; AND
 - iii. Patient has responded to Benlysta subcutaneous or intravenous, as determined by the prescriber.
Note: Examples of a response include improvement in organ dysfunction, reduction in flares, reduction in corticosteroid dose, decrease of anti-dsDNA titer, and improvement in complement levels (i.e., C3, C4).
2. **Systemic Lupus Erythematosus.** Approve for the duration noted if the patient meets ONE of the following (A or B):
 - A) **Initial Therapy.** Approve for 4 months if the patient meets ALL of the following (i, ii, iii, and iv):
 - i. Patient is ≥ 5 years of age; AND
 - ii. Patient has autoantibody-positive systemic lupus erythematosus (SLE), defined as positive for antinuclear antibodies (ANA) and/or anti-double-stranded DNA (anti-dsDNA) antibody; AND
Note: Not all patients with SLE are positive for anti-dsDNA, but most will be positive for ANA.
 - iii. Patient meets ONE of the following (a or b):
 - a) The medication is being used concurrently with at least one other standard therapy; OR
Note: Examples of standard therapies include an antimalarial (e.g., hydroxychloroquine), systemic corticosteroid (e.g., prednisone), and other immunosuppressants (e.g., azathioprine, mycophenolate mofetil, methotrexate).

- b) Patient is determined to be intolerant to standard therapy due to a significant toxicity, as determined by the prescriber; AND
 - iv. The medication is prescribed by or in consultation with a rheumatologist, clinical immunologist, nephrologist, neurologist, or dermatologist; OR
- B) Patient is Currently Receiving Benlysta Intravenous or Subcutaneous.** Approve for 1 year if the patient meets ALL of the following (i, ii, and iii):
- i. Patient meets ONE of the following (a or b):
 - a) The medication is being used concurrently with at least one other standard therapy; OR
Note: Examples of standard therapies include an antimalarial (e.g., hydroxychloroquine), systemic corticosteroid (e.g., prednisone), and other immunosuppressants (e.g., azathioprine, mycophenolate mofetil, methotrexate).
 - b) Patient is determined to be intolerant to standard therapy due to a significant toxicity, as determined by the prescriber; AND
 - ii. The medication is prescribed by or in consultation with a rheumatologist, clinical immunologist, nephrologist, neurologist, or dermatologist; AND
 - iii. Patient has responded to Benlysta subcutaneous or intravenous, as determined by the prescriber.
Note: Examples of a response include reduction in flares, reduction in corticosteroid dose, decrease of anti-dsDNA titer, improvement in complement levels (i.e., C3, C4), or improvement in specific organ dysfunction (e.g., musculoskeletal, blood, hematologic, vascular, others).

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Benlysta intravenous is not recommended in the following situations:

- 1. Concurrent Use with Other Biologics.** Benlysta intravenous has not been studied and is not recommended in combination with other biologics.¹ Safety and efficacy have not been established with these combinations. See [APPENDIX](#) for examples of biologics that should not be taken in combination with Benlysta.
- 2. Concurrent Use with Lupkynis (voclosporin capsules).** Lupkynis has not been studied in combination with biologics such as Benlysta.¹
- 3. Rheumatoid Arthritis.** A Phase II dose-ranging study evaluating patients with rheumatoid arthritis showed only small American College of Rheumatology (ACR) 20 responses with Benlysta (e.g., ACR 20 response at Week 24 was 28% with Benlysta 10 mg/kg).⁴ Numerous other agents are available with higher ACR responses and established efficacy for RA.
- 4.** Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Benlysta® injection [prescribing information]. Durham, NC: GlaxoSmithKline; June 2024.
2. Fanouriakis A, Kostopoulou M, Andersen J, et al. EULAR recommendations for the management of systemic lupus erythematosus: 2023 update. *Ann Rheum Dis.* 2024;83(1):15-29.
3. Kidney Disease: Improving Global Outcomes (KDIGO) Lupus Nephritis Work Group. KDIGO 2024 Clinical Practice Guideline for the management of LUPUS NEPHRITIS. *Kidney Int.* 2024;105(1S):S1-S69.
4. Stohl W, Merrill JT, McKay JD, et al. Efficacy and safety of belimumab in patients with rheumatoid arthritis: a phase II, randomized, double-blind, placebo-controlled, dose-ranging study. *J Rheumatol.* 2013;40(5):579-589.

HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	No criteria changes.	03/08/2023
Selected Revision	Lupus Nephritis: For initial therapy, a requirement was added that the patient has biopsy-confirmed lupus nephritis. For initial therapy and a patient currently receiving Benlysta, the requirement that the patient is taking with standard therapy was changed to more generally require that the patient is taking an immunosuppressive regimen. Leflunomide, methotrexate, and/or systemic corticosteroids were added to existing concurrent medication examples. The exception for a patient who is intolerant to standard therapy due to significant toxicity as determined by the prescriber was removed from the Policy.	04/26/2023
Selected Revision	Lupus Nephritis: For initial therapy, the requirement that the “Patient has autoantibody-positive systemic lupus erythematosus (SLE), defined as positive for antinuclear antibodies (ANA) and/or anti-double-stranded DNA (anti-dsDNA) antibody” was removed from the Policy.	07/05/2023
Annual Revision	No criteria changes.	03/13/2024
Update	06/04/2024: No criteria changes. Overview: The indication for systemic lupus erythematosus was updated to remove “auto-antibody positive” and following statement was added, “The efficacy of Benlysta for the treatment of SLE was studied in patients with a history of autoantibodies (anti-nuclear antibody and/or anti-double-stranded DNA) and an exploratory analysis of the pivotal trial indicated Benlysta was beneficial in patients who were autoantibody positive.” Additionally, the following statement was removed due to the labeling change for Benlysta subcutaneous for systemic lupus erythematosus, “Of note, there is also a subcutaneous formulation of Benlysta with a similar indication except use is limited to those $\geq$ 18 years of age.”	--
Annual Revision	No criteria changes. Updated Appendix.	03/19/2025

APPENDIX

	Mechanism of Action	Examples of Indications*
Biologics		
Saphnelo® (anifrolumab-fnia IV infusion)	IFN receptor antagonist	SLE
Adalimumab SC Products (Humira®, biosimilars)	Inhibition of TNF	AS, CD, JIA, PsO, PsA, RA, UC
Cimzia® (certolizumab pegol SC injection)	Inhibition of TNF	AS, CD, nr-axSpA, PsO, PsA, RA
Etanercept SC Products (Enbrel®, biosimilars)	Inhibition of TNF	AS, JIA, PsO, PsA, RA
Infliximab IV Products (Remicade®, biosimilars)	Inhibition of TNF	AS, CD, PsO, PsA, RA, UC
Zymfentra® (infliximab-dyyb SC injection)	Inhibition of TNF	CD, UC
Simponi®, Simponi Aria® (golimumab SC injection, golimumab IV infusion)	Inhibition of TNF	SC formulation: AS, PsA, RA, UC
		IV formulation: AS, PJIA, PsA, RA
Tocilizumab Products (Actemra® IV, biosimilar; Actemra SC, biosimilar)	Inhibition of IL-6	SC formulation: PJIA, RA, SJIA
		IV formulation: PJIA, RA, SJIA
Kevzara® (sarilumab SC injection)	Inhibition of IL-6	RA
Orencia® (abatacept IV infusion, abatacept SC injection)	T-cell costimulation modulator	SC formulation: JIA, PSA, RA
		IV formulation: JIA, PsA, RA
Rituximab IV Products (Rituxan®, biosimilars)	CD20-directed cytolytic antibody	RA
Kineret® (anakinra SC injection)	Inhibition of IL-1	JIA [^] , RA
Omvo® (mirikizumab IV infusion, SC injection)	Inhibition of IL-23	UC
Stelara® (ustekinumab SC injection, ustekinumab IV infusion)	Inhibition of IL-12/23	SC formulation: CD, PsO, PsA, UC
		IV formulation: CD, UC
Siliq® (brodalumab SC injection)	Inhibition of IL-17	PsO
Cosentyx® (secukinumab SC injection; secukinumab IV infusion)	Inhibition of IL-17A	SC formulation: AS, ERA, nr-axSpA, PsO, PsA
		IV formulation: AS, nr-axSpA, PsA
Taltz® (ixekizumab SC injection)	Inhibition of IL-17A	AS, nr-axSpA, PsO, PsA
Bimzelx® (bimekizumab-bkzx SC injection)	Inhibition of IL-17A/17F	PsO
Ilumya® (tildrakizumab-asmn SC injection)	Inhibition of IL-23	PsO
Skyrizi® (risankizumab-rzaa SC injection, risankizumab-rzaa IV infusion)	Inhibition of IL-23	SC formulation: CD, PSA, PsO, UC
		IV formulation: CD, UC
Tremfya® (guselkumab SC injection, guselkumab IV infusion)	Inhibition of IL-23	SC formulation: PsA, PsO, UC
		IV formulation: UC
Entyvio® (vedolizumab IV infusion, vedolizumab SC injection)	Integrin receptor antagonist	CD, UC

* Not an all-inclusive list of indication (e.g., oncology indications and rare inflammatory conditions are not listed). Refer to the prescribing information for the respective agent for FDA-approved indications; SC – Subcutaneous; IV – Intravenous; BLyS – B-lymphocyte stimulator-specific inhibitor; SLE – Systemic lupus erythematosus; IFN – Interferon; TNF – Tumor necrosis factor; AS – Ankylosing spondylitis; CD – Crohn’s disease; JIA – Juvenile idiopathic arthritis; PsO – Plaque psoriasis; PsA – Psoriatic arthritis; RA – Rheumatoid arthritis; UC – Ulcerative colitis; nr-axSpA – Non-radiographic axial spondyloarthritis; PJIA – Polyarticular juvenile idiopathic arthritis; IL – Interleukin; SJIA – Systemic juvenile idiopathic arthritis; [^] Off-label use of Kineret in JIA supported in guidelines; ERA – Enthesitis-related arthritis.