

CARE VALUE POLICY

POLICY: Calcitonin Gene-Related Peptide Inhibitors – Emgality Care Value Policy

- Emgality® (galcanezumab-gnlm subcutaneous injection – Lilly)

REVIEW DATE: 04/21/2021; selected revision 07/21/2021

OVERVIEW

Emgality, a calcitonin gene-related peptide (CGRP) antagonist, is indicated for the following uses:¹

- **Episodic cluster headache treatment** in adults.
- **Migraine headache prevention** in adults.

Disease Overview

Migraines have been defined as chronic or episodic. Chronic migraine is described by the International Headache Society as headache occurring on ≥ 15 days/month for > 3 months and has the features of migraine headache on ≥ 8 days/month.² Episodic migraine is characterized by headaches that occur < 15 days/month.^{3,4} Episodic migraine is more common than chronic migraine; however, chronic migraine is associated with a markedly greater personal and societal burden.

Cluster headaches are associated with attacks of severe, strictly unilateral pain which is orbital, supraorbital, temporal, or in any combination of these sites, lasting 15 to 180 minutes.² The headaches occur from once every other day to eight times a day. Cluster headache is considered among the most severe of the primary headache disorders because of extreme pain, associated autonomic symptoms, and high attack frequency.⁵ In addition, a large proportion of patients with cluster headache have chronic cluster headache, which features only brief or no remission periods, and may be particularly refractory to medical therapies.

Guidelines

An updated assessment of the **preventive and acute treatment of migraine** by the **American Headache Society (AHS)** [2018] reaffirms previous migraine guidelines.⁶ Patients with migraine should be considered for preventive treatment when attacks significantly interfere with patients' daily routines despite acute treatment; frequent attacks (≥ 4 monthly headache days); contraindication to, failure, overuse, or adverse events with acute treatments; or patient preference. Before developing a preventive treatment plan, the appropriate use (e.g., drug type, route and timing of administration, frequency) of acute treatments should be initiated and coupled with education and lifestyle modifications. All patients with migraine should be offered a trial of acute treatment. Based on the level of evidence for efficacy and the American Academy of Neurology scheme for classification of evidence, the following oral treatments have established efficacy and should be offered for migraine prevention: antiepileptic drugs (divalproex sodium, valproate sodium, topiramate [not for women of childbearing potential without a reliable method of birth control]); beta-blockers (metoprolol, propranolol, timolol); and frovatriptan (for short-term preventive treatment of menstrual migraine).⁷ The following treatments are probably effective and should be considered for migraine prevention: antidepressants (amitriptyline, venlafaxine); beta-blockers (atenolol, nadolol); and angiotensin receptor blockers (candesartan).

Four injectable preventive therapies for migraine are mentioned in the AHS consensus statement: Botox® (onabotulinumtoxinA injection) and three monoclonal antibodies targeting CGRP (Aimovig® [erenumab-aooe injection], Ajovy® [fremanezumab-vfrm injection], and Emgality).⁶ Of note, Vyepti™ (eptinezumab-jjmr injection) had not been approved at the time of the consensus statement. The update notes that a CGRP inhibitor should only be initiated in patients who are diagnosed with migraine, have ≥ 4 migraine headache days per month, and have intolerance or inadequate response to 6-week trials of at least two traditional oral

migraine preventive medications. Additional criteria apply depending on the number and severity of monthly headache days. Clinical judgment may result in an emerging treatment being added to one or more established treatments. If initiating treatment with a CGRP inhibitor in a patient already on a preventive treatment, it is appropriate to add the CGRP inhibitor to the existing regimen and make no other changes until the effectiveness of the CGRP inhibitor is determined since the risk of interactions between traditional oral migraine preventive medications and the CGRP inhibitors is minimal or nonexistent. Making a decision regarding continuation of therapy for a CGRP inhibitor requires a trial of the medication for at least 3 months, and treatment should be continued only if benefits can be documented during that time (e.g., reduction in mean monthly headache days of $\geq 50\%$ relative to the pretreatment baseline). Since migraine may improve or remit over time, it is important to reevaluate therapeutic response and, if possible, taper or discontinue treatment if patients no longer meet the criteria for offering preventive treatment. However, once control is established, the decision to discontinue or taper treatment should be a shared decision between patient and clinician.

The AHS has published evidence-based guidelines on the **treatment of cluster headache** (2016).⁵ The guidelines recommend sumatriptan subcutaneous, zolmitriptan nasal spray, and high flow oxygen for acute treatment. For prophylactic therapy, suboccipital steroid injection has been established as effective for the prophylactic therapy of episodic and chronic cluster headache (Level A). Lithium, verapamil, and melatonin are considered possibly effective for the prophylactic therapy of episodic and chronic cluster headache (Level C). Currently, there is insufficient evidence to make a recommendation for frovatriptan and prednisone (Level U).

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Emgality. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Emgality is recommended in those who meet the following criteria:

FDA-Approved Indications

- 1. Episodic Cluster Headache Treatment.** Approve Emgality for 6 months if the patient meets the following criteria (A, B, C, and D):
 - A)** Patient is ≥ 18 years of age; AND
 - B)** Patient has between one headache every other day and eight headaches per day; AND
 - C)** Patient has tried at least one standard prophylactic (preventive) pharmacologic therapy for cluster headache; AND
Note: Examples of standard prophylactic (preventive) pharmacologic therapies for cluster headache include lithium, verapamil, melatonin, frovatriptan, prednisone, suboccipital steroid injection, topiramate, and valproate.
 - D)** Patient has had inadequate efficacy or has experienced adverse event(s) severe enough to warrant discontinuation of the standard prophylactic (preventive) pharmacologic therapy, according to the prescriber.
-

2. Migraine Headache Prevention. Approve Emgality for 1 year if the patient meets the following criteria (A, B, C, D and E):

A) Patient is ≥ 18 years of age; AND

B) Patient has ≥ 4 migraine headache days per month (prior to initiating a migraine-preventative medication); AND

C) Patient has tried at least two standard prophylactic (preventive) pharmacologic therapies, each from a different pharmacologic class; AND

Note: Examples of standard prophylactic (preventive) pharmacologic therapies for migraine include angiotensin receptor blocker, angiotensin converting enzyme inhibitor, anticonvulsant, beta-blocker, calcium channel blocker, tricyclic antidepressant, other antidepressant.

D) Patient meets ONE of the following criteria (i, ii, or iii):

i. Patient has had inadequate efficacy to both of those standard prophylactic (preventive) pharmacologic therapies, according to the prescriber; OR

ii. Patient has experienced adverse event(s) severe enough to warrant discontinuation of both of those standard prophylactic (preventive) pharmacologic therapies, according to the prescriber; OR

iii. Patient has had inadequate efficacy to one standard prophylactic (preventive) pharmacologic therapy and has experienced adverse event(s) severe enough to warrant discontinuation to another standard prophylactic (preventive) pharmacologic therapy, according to the prescriber; AND

E) Patient meets ONE of the following (i, ii, or iii):

i. Patient is NOT taking a calcitonin gene-related peptide (CGRP) inhibitor for migraine headache prevention and meets ONE of the following (a or b):

Note: CGRP inhibitors used for migraine headache prevention are Aimovig (erenumab-aooe injection), Ajovy (fremanezumab-vfrm injection), Emgality, and Vyepti (eptinezumab-jjmr injection).

a) Patient has tried at least one triptan therapy; OR

b) Patient has a contraindication to triptan(s) according to the prescriber; OR

Note: Examples of contraindications to triptans include a history of coronary artery disease; cardiac accessory conduction pathway disorders; history of stroke, transient ischemic attack, or hemiplegic or basilar migraine; peripheral vascular disease; ischemic bowel disease; uncontrolled hypertension; or severe hepatic impairment.

ii. Patient is currently taking Emgality and has had a significant clinical benefit from the medication as determined by the prescriber; OR

Note: Examples of significant clinical benefit include a reduction in the overall number of migraine days per month or a reduction in number of severe migraine days per month from the time that Emgality was initiated.

iii. Patient is switching from a different CGRP inhibitor for migraine headache prevention to Emgality.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Emgality is not recommended in the following situations:

1. **Acute Treatment of Migraine.** Emgality has not been studied for the acute treatment of migraine.
2. **Combination Therapy with Aimovig (erenumab-aooe injection for subcutaneous use), Ajovy (fremanezumab-vfrm injection for subcutaneous use), Vyepti (eptinezumab-jjmr injection for intravenous use), or preventive treatment of migraine with Nurtec™ ODT (rimegepant sulfate orally disintegrating tablet).** Ajovy, Aimovig, Emgality, and Vyepti are calcitonin gene-related peptide (CGRP) inhibitors for migraine prevention and have not been studied for use in combination with another agent in the same class.⁸⁻¹⁰ Nurtec ODT is an oral CGRP inhibitor for the acute treatment of migraine and for the preventive treatment of episodic migraine in adults.¹¹
3. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Emgality® injection for subcutaneous use [prescribing information]. Indianapolis, IN: Eli Lilly and Company; December 2019.
 2. Headache Classification Subcommittee of the International Headache Society. The International Classification of Headache Disorders: 3rd edition (beta version). *Cephalalgia*. 2013;33:629-808.
 3. MacGregor EA. In the clinic. Migraine. *Ann Int Med*. 2013;159:ITC5-1-ITC5-16.
 4. Lipton RB, Silberstein SD. Episodic and chronic migraine headache: breaking down barriers to optimal treatment and prevention. *Headache*. 2015;52:103-122.
 5. Robbins MS, Starling AJ, Pringsheim TM, et al. Treatment of cluster headache: the American Headache Society evidence-based guidelines. *Headache*. 2016;56:1093-1106.
 6. American Headache Society. The American Headache Society position statement on integrating new migraine treatments into clinical practice. *Headache*. 2019;59:1-18.
 7. Silberstein SD, Holland S, Freitag F, et al. Evidence-based guideline update: Pharmacologic treatment for episodic migraine prevention in adults. Report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Headache Society. *Neurology*. 2012;78(17):1337-1345.
 8. Aimovig® injection for subcutaneous use [prescribing information]. Thousand Oaks, CA: Amgen; February 2021.
 9. Ajovy® injection for subcutaneous use [prescribing information]. North Wales, PA: Teva Pharmaceuticals USA, Inc.; January 2020.
 10. Vyepti™ injection for intravenous use [prescribing information]. Bothell, WA: Lundbeck Seattle BioPharmaceuticals, Inc.; February 2020.
 11. Nurtec™ ODT [prescribing information]. New Haven, CT: Biohaven Pharmaceuticals; May 2021.
-

HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	Condition Not Recommended for Approval of Combination Therapy: Vyepti was added to the list of injectable calcitonin gene-related peptide inhibitors.	04/15/2020
Annual Revision	Migraine Headache Prevention: Added criterion for patients who are already taking a calcitonin gene-related peptide (CGRP) inhibitor requiring the patient to have had a significant clinical benefit from the medication as determined by the prescriber. Changed the criteria asking about previous triptan use to only apply to patients who are not taking a CGRP inhibitor. Moved examples of standard prophylactic (preventive) pharmacologic therapies into a note. Added “(preventive)” following the word “prophylactic” as clarification. Episodic Cluster Headache Treatment: Added “(preventive)” following the word “prophylactic” as clarification.	04/21/2021
Selected Revision	Migraine Headache Prevention: The criterion “Patient is currently taking a CGRP inhibitor for migraine headache prevention” was changed to “Patient is currently taking Emgality” and an additional criterion was added for “Patient is switching from a different CGRP inhibitor for migraine headache prevention to Emgality”. Also, a note was added to provide examples of contraindications to triptan use. Conditions Not Recommended for Approval: Preventive treatment of migraine with Nurtec ODT was added to the list of injectable CGRP inhibitors that cannot be used in combination with Emgality.	07/21/2021